

CODEX ALIMENTARIUS

INTERNATIONAL FOOD STANDARDS



Food and Agriculture
Organization of
the United Nations



World Health
Organization

E-mail: codex@fao.org - www.codexalimentarius.org

MAXIMUM RESIDUE LIMITS (MRLs) AND RISK MANAGEMENT RECOMMENDATIONS (RMRs) FOR RESIDUES OF VETERINARY DRUGS IN FOODS

CAC/MRL 2-2015

Updated as at the 38th Session of the Codex Alimentarius Commission (July 2015)

Maximum Residue Limits (MRL)

Abamectin	Flumequine
Albendazole	Gentamicin
Amoxicillin	Imidocarb
Avylamycin	Isometamidium
Azaperone	Ivermectin
Benzylpenicillin/Procaine benzylpenicillin	Levamisole
Carazolol	Lincomycin
Ceftiofur	Melengestrol acetate
Chlortetracycline/Oxytetracycline/Tetracycline	Monensin
Clenbuterol	Monepantel
Closantel	Moxidectin
Colistin	Narasin
Cyfluthrin	Neomycin
Cyhalothrin	Nicarbazin
Cypermethrin and alpha-cypermethrin	Phoxim
Danofloxacin	Pirlimycin
Deltamethrin	Porcine somatotropin
Derquantel	Progesterone
Dexamethasone	Ractopamine
Diclazuril	Sarafloxacin
Dicyclanil	Spectinomycin
Dihydrostreptomycin/Streptomycin	Spiramycin
Diminazene	Sulfadimidine
Doramectin	Testosterone
Enamectin benzoate	Thiabendazole
Eprinomectin	Tilmicosin
Erythromycin	Trenbolone acetate
Estradiol-17beta	Trichlorfon (Metrifonate)
Febantel/Fenbendazole/Oxfendazole	Triclabendazole
Fluazuron	Tylosin
Flubendazole	Zeranol

Risk Management Recommendations (RMR) for Residues of Veterinary Drugs

Carbadox	Metronidazole
Chloramphenicol	Nitrofur
Chlorpromazine	Olaquinox
Dimetridazole	Ronidazole
Furazolidone	Stilbens
Ipronidazole	
Malachite Green	

MAXIMUM RESIDUE LIMITS (MRLs) FOR RESIDUES OF VETERINARY DRUGS IN FOODS

ABAMECTIN (anthelmintic agent)				
JECFA Evaluation: 45 (1995); 47 (1996)				
Acceptable Daily Intake : 0-2 µg/kg body weight (1997) Established for the sum of abamectin and (Z)-8,9 isomer by the 1997 JMPR.				
Residue Definition: Avermectin B1a.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Liver	100	26 th (2003)	
Cattle	Kidney	50	26 th (2003)	
Cattle	Fat	100	26 th (2003)	

ALBENDAZOLE (anthelmintic agent)				
JECFA Evaluation: 34 (1989)				
Acceptable Daily Intake: 0-50 µg/kg body weight (34 th JECFA, 1989).				
Residue Definition: Except milk, 2-aminosulfone metabolite; Milk, not yet identified.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Not specified	Muscle	100	20 th (1993)	
Not specified	Liver	5000	20 th (1993)	
Not specified	Kidney	5000	20 th (1993)	
Not specified	Fat	100	20 th (1993)	
Not specified	Milk (µg/l)	100	20 th (1993)	

AMOXICILLIN (antimicrobial agent)**JECFA Evaluation:** 75 (2011)**Acceptable Daily Intake:** 0-0.07 µg/kg body weight on the basis of microbiological effects (75th JECFA, 2011).**Estimated Dietary Exposure** The 75th JECFA (2001) did not calculate an EDI for amoxicillin owing to the small number of quantifiable residue data points. Using the model diet of 300 g muscle, 100 g live, 50 g kidney, 50 g fat and 1.5 liter of milk with the MRLs recommended, the theoretical maximum daily intake (TMDI) is 31 µg/person, which represents 74% of the upper bound of the ADI.**Residue Definition:** Amoxicillin

Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	50	35 th (2012)	
Cattle	Liver	50	35 th (2012)	
Cattle	Kidney	50	35 th (2012)	
Cattle	Fat	50	35 th (2012)	
Cattle	Milk	4	35 th (2012)	
Sheep	Muscle	50	35 th (2012)	
Sheep	Liver	50	35 th (2012)	
Sheep	Kidney	50	35 th (2012)	
Sheep	Fat	50	35 th (2012)	
Sheep	Milk	4	35 th (2012)	
Pigs	Muscle	50	35 th (2012)	
Pigs	Liver	50	35 th (2012)	
Pigs	Kidney	50	35 th (2012)	
Pigs	Fat/Skin	50	35 th (2012)	

AVILAMYCIN (antimicrobial agent)				
JECFA Evaluation: 70 (2008)				
Acceptable Daily Intake: 0-2 mg/kg body weight on the basis of a NOAEL of 150 mg avilamycin activity/kg body weight per day and a safety factor of 100 and rounding to one significant figure (70 th JECFA, 2008).				
Residue Definition: Dichloroisoeverninic acid (DIA).				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Pigs	Muscle	200	32 nd (2009)	
Pigs	Liver	300	32 nd (2009)	
Pigs	Kidney	200	32 nd (2009)	
Pigs	Fat/Skin	200	32 nd (2009)	
Chicken	Muscle	200	32 nd (2009)	
Chicken	Liver	300	32 nd (2009)	
Chicken	Kidney	200	32 nd (2009)	
Chicken	Fat/Skin	200	32 nd (2009)	
Turkey	Muscle	200	32 nd (2009)	
Turkey	Liver	300	32 nd (2009)	
Turkey	Kidney	200	32 nd (2009)	
Turkey	Fat/Skin	200	32 nd (2009)	
Rabbits	Muscle	200	32 nd (2009)	
Rabbits	Liver	300	32 nd (2009)	
Rabbits	Kidney	200	32 nd (2009)	
Rabbits	Fat/Skin	200	32 nd (2009)	

AZAPERONE (tranquilizing agent)				
JECFA Evaluation: 38 (1991); 43 (1994); 50 (1998); 52 (1999)				
Acceptable Daily Intake: 0-6 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: Sum of azaperone and azaperol.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Pig	Muscle	60	23 rd (1999)	
Pig	Liver	100	23 rd (1999)	
Pig	Kidney	100	23 rd (1999)	
Pig	Fat	60	23 rd (1999)	

BENZYL PENICILLIN/PROCAINE BENZYL PENICILLIN (antimicrobial agent)				
JECFA Evaluation: 36 (1990); 50 (1998)				
Acceptable Daily Intake: 30 µg-penicillin/person/day (50 th JECFA, 1998). Residues of benzylpenicillin and procaine benzylpenicillin should be kept below this level.				
Residue Definition: Benzylpenicillin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	50	23 rd (1999)	
Cattle	Liver	50	23 rd (1999)	
Cattle	Kidney	50	23 rd (1999)	
Cattle	Milk (µg/l)	4	23 rd (1999)	
Chicken	Muscle	50	23 rd (1999)	Applies to procaine benzylpenicillin only.
Chicken	Liver	50	23 rd (1999)	Applies to procaine benzylpenicillin only.
Chicken	Kidney	50	23 rd (1999)	Applies to procaine benzylpenicillin only.
Pig	Muscle	50	23 rd (1999)	
Pig	Liver	50	23 rd (1999)	
Pig	Kidney	50	23 rd (1999)	

CARAZOLOL (beta-adrenoceptor blocking agent)				
JECFA Evaluation: 38 (1991); 43 (1994); 52 (1999)				
Acceptable Daily Intake: 0-0.1 µg/kg body weight (43 rd JECFA, 1994). ADI based on the acute pharmacological effects of carazolol.				
Residue Definition: Carazolol.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Pig	Muscle	5	26 th (2003)	The concentration at the injection site two hours after treatment may result in an intake that exceeds the acute RfD and therefore, an appropriate withdrawal period should be applied.
Pig	Liver	25	26 th (2003)	
Pig	Kidney	25	26 th (2003)	
Pig	Fat/Skin	5	26 th (2003)	The concentration at the injection site two hours after treatment may result in an intake that exceeds the acute RfD and therefore, an appropriate withdrawal period should be applied.

CEFTIOFUR (antimicrobial agent)				
JECFA Evaluation: 45 (1995); 48 (1997)				
Acceptable Daily Intake: 0-50 µg/kg body weight (45 th JECFA, 1995).				
Residue Definition: Desfuroylceftiofur.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	1000	23 rd (1999)	
Cattle	Liver	2000	23 rd (1999)	
Cattle	Kidney	6000	23 rd (1999)	
Cattle	Fat	2000	23 rd (1999)	
Cattle	Milk (µg/l)	100	23 rd (1999)	
Pig	Muscle	1000	23 rd (1999)	
Pig	Liver	2000	23 rd (1999)	
Pig	Kidney	6000	23 rd (1999)	
Pig	Fat	2000	23 rd (1999)	

CHLORTETRACYCLINE/OXYTETRACYCLINE/TETRACYCLINE (antimicrobial agent)**JECFA Evaluation:** 45 (1995); 47 (1996); 50 (1998); 58 (2002)**Acceptable Daily Intake:** 0-30 µg/kg body weight (50th JECFA, 1998). Group ADI for chlortetracycline, oxytetracycline and tetracycline.**Residue Definition:** Parent drugs, singly or in combination.

Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	200	26 th (2003)	
Cattle	Liver	600	26 th (2003)	
Cattle	Kidney	1200	26 th (2003)	
Cattle	Milk (µg/l)	100	26 th (2003)	
Fish	Muscle	200	26 th (2003)	Applies only to oxytetracycline.
Giant prawn (<i>Paeneus monodon</i>)	Muscle	200	26 th (2003)	Applies only to oxytetracycline.
Pig	Muscle	200	26 th (2003)	
Pig	Liver	600	26 th (2003)	
Pig	Kidney	1200	26 th (2003)	
Poultry	Muscle	200	26 th (2003)	
Poultry	Liver	600	26 th (2003)	
Poultry	Kidney	1200	26 th (2003)	
Poultry	Eggs	400	26 th (2003)	
Sheep	Muscle	200	26 th (2003)	
Sheep	Liver	600	26 th (2003)	
Sheep	Kidney	1200	26 th (2003)	
Sheep	Milk (µg/l)	100	26 th (2003)	

CLENBUTEROL (adrenoceptor agonist)				
JECFA Evaluation: 47 (1996)				
Acceptable Daily Intake: 0-0.004 µg/kg body weight (47 th JECFA, 1996).				
Residue Definition: Clenbuterol.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	0.2	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Liver	0.6	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Kidney	0.6	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Fat	0.2	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Milk (µg/l)	0.05	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Muscle	0.2	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Liver	0.6	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Kidney	0.6	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Fat	0.2	26 th (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.

CLOSANTEL (anthelmintic agent)				
JECFA Evaluation: 36 (1990); 40 (1992)				
Acceptable Daily Intake: 0-30 µg/kg body weight (40 th JECFA, 1992).				
Residue Definition: Closantel.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	1000	20 th (1993)	
Cattle	Liver	1000	20 th (1993)	
Cattle	Kidney	3000	20 th (1993)	
Cattle	Fat	3000	20 th (1993)	
Sheep	Muscle	1500	20 th (1993)	
Sheep	Liver	1500	20 th (1993)	
Sheep	Kidney	5000	20 th (1993)	
Sheep	Fat	2000	20 th (1993)	

COLISTIN (antimicrobial agent)				
JECFA Evaluation: 66 (2006)				
Acceptable Daily Intake: 0-7 µg/kg body weight (66 th JECFA, 2006).				
Residue Definition: Sum of colistin A and colistin B.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	150	31 st (2008)	
Cattle	Liver	150	31 st (2008)	
Cattle	Kidney	200	31 st (2008)	
Cattle	Fat	150	31 st (2008)	
Cattle	Milk	50	31 st (2008)	
Sheep	Muscle	150	31 st (2008)	
Sheep	Liver	150	31 st (2008)	
Sheep	Kidney	200	31 st (2008)	
Sheep	Fat	150	31 st (2008)	
Sheep	Milk	50	31 st (2008)	
Goat	Muscle	150	31 st (2008)	
Goat	Liver	150	31 st (2008)	
Goat	Kidney	200	31 st (2008)	
Goat	Fat	150	31 st (2008)	
Pig	Muscle	150	31 st (2008)	
Pig	Liver	150	31 st (2008)	
Pig	Kidney	200	31 st (2008)	
Pig	Fat	150	31 st (2008)	The MRL includes skin + fat
Chicken	Muscle	150	31 st (2008)	
Chicken	Liver	150	31 st (2008)	
Chicken	Kidney	200	31 st (2008)	
Chicken	Fat	150	31 st (2008)	The MRL includes skin + fat
Chicken	Eggs	300	31 st (2008)	
Turkey	Muscle	150	31 st (2008)	
Turkey	Liver	150	31 st (2008)	
Turkey	Kidney	200	31 st (2008)	
Turkey	Fat	150	31 st (2008)	The MRL includes skin + fat
Rabbit	Muscle	150	31 st (2008)	
Rabbit	Liver	150	31 st (2008)	
Rabbit	Kidney	200	31 st (2008)	
Rabbit	Fat	150	31 st (2008)	

CYFLUTHRIN (insecticide)				
JECFA Evaluation: 48 (1997)				
Acceptable Daily Intake: 0-20 µg/kg body weight (48 th JECFA, 1997).				
Residue Definition: Cyfluthrin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	20	26 th (2003)	
Cattle	Liver	20	26 th (2003)	
Cattle	Kidney	20	26 th (2003)	
Cattle	Fat	200	26 th (2003)	
Cattle	Milk (µg/l)	40	26 th (2003)	

CYHALOTHRIN (insecticide)				
JECFA Evaluation: 54 (2000); 58 (2002); 62 (2004)				
Acceptable Daily Intake: 0-5 µg/kg body weight (62 nd JECFA, 2004).				
Residue Definition: Cyhalothrin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	20	28 th (2005)	
Cattle	Liver	20	28 th (2005)	
Cattle	Kidney	20	28 th (2005)	
Cattle	Fat	400	28 th (2005)	
Cattle	Milk	30	28 th (2005)	
Pig	Muscle	20	28 th (2005)	
Pig	Liver	20	28 th (2005)	
Pig	Kidney	20	28 th (2005)	
Pig	Fat	400	28 th (2005)	
Sheep	Muscle	20	28 th (2005)	
Sheep	Liver	50	28 th (2005)	
Sheep	Kidney	20	28 th (2005)	
Sheep	Fat	400	28 th (2005)	

CYPERMETHRIN AND ALPHA-CYPERMETHRIN (insecticide)				
JECFA Evaluation: 62 (2004)				
Acceptable Daily Intake: JECFA established a common ADI of 0-20 µg/kg bw for both cypermethrin and alpha-cypermethrin (62 nd JECFA, 2004)..				
Residue Definition: Total of cypermethrin residues (resulting from the use of cypermethrin or alpha-cypermethrin as veterinary drugs).				
Species	Tissue	MRLs(µg/kg)	CAC	Note
Cattle	Muscle	50	29 th (2006)	
Cattle	Liver	50	29 th (2006)	
Cattle	Kidney	50	29 th (2006)	
Cattle	Fat	1000	29 th (2006)	
Cattle	Milk	100	29 th (2006)	
Sheep	Muscle	50	29 th (2006)	
Sheep	Liver	50	29 th (2006)	
Sheep	Kidney	50	29 th (2006)	
Sheep	Fat	1000	29 th (2006)	

DANOFLOXACIN (antimicrobial agent)				
JECFA Evaluation: 48 (1997)				
Acceptable Daily Intake: 0-20 µg/kg body weight (48 th JECFA, 1997).				
Residue Definition: Danofloxacin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	200	24 th (2001)	
Cattle	Liver	400	24 th (2001)	
Cattle	Kidney	400	24 th (2001)	
Cattle	Fat	100	24 th (2001)	
Chicken	Muscle	200	24 th (2001)	
Chicken	Liver	400	24 th (2001)	
Chicken	Kidney	400	24 th (2001)	
Chicken	Fat	100	24 th (2001)	Fat/skin in normal proportion.
Pig	Muscle	100	24 th (2001)	
Pig	Liver	50	24 th (2001)	
Pig	Kidney	200	24 th (2001)	
Pig	Fat	100	24 th (2001)	

DELTAMETHRIN (insecticide)				
JECFA Evaluation: 52 (1999); 60 (2003)				
Acceptable Daily Intake: 0-10 µg/kg body weight (1982). Established by the 1982 JMPR.				
Residue Definition: Deltamethrin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	30	26 th (2003)	
Cattle	Liver	50	26 th (2003)	
Cattle	Kidney	50	26 th (2003)	
Cattle	Fat	500	26 th (2003)	
Cattle	Milk	30	26 th (2003)	
Chicken	Muscle	30	26 th (2003)	
Chicken	Liver	50	26 th (2003)	
Chicken	Kidney	50	26 th (2003)	
Chicken	Fat	500	26 th (2003)	
Chicken	Eggs	30	26 th (2003)	
Salmon	Muscle	30	26 th (2003)	
Sheep	Muscle	30	26 th (2003)	
Sheep	Liver	50	26 th (2003)	
Sheep	Kidney	50	26 th (2003)	
Sheep	Fat	500	26 th (2003)	

DERQUANTEL (anthelmintic agent)				
JECFA Evaluation: 75 (2011); 78 (2013)				
Acceptable Daily Intake: 0-0.3 µg/kg body weight on the basis of a lowest-observed-adverse-effect level (LOAEL) of 0.1 mg/kg body weight per day for acute clinical observations in dogs, consistent with antagonistic activity on the nicotinic acetylcholine receptors. A safety factor of 300 was applied to the LOAEL (75 th JECFA, 2011).				
Estimated Dietary Exposure: There were insufficient data to calculate an EDI, and the TMDI approach was used. Using the model diet and the MT:TR approach, these MRLs result in an estimated dietary exposure of 6.8 µg/person, which represents approximately 38% of the upper bound of the ADI (78 th JECFA, 2013).				
Residue Definition: Derquantel.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Sheep	Muscle	0.3	38 th (2015)	
Sheep	Liver	0.8	38 th (2015)	
Sheep	Kidney	0.4	38 th (2015)	
Sheep	Fat	7.0	38 th (2015)	

DEXAMETHASONE (glucocorticosteroid)				
JECFA Evaluation: 70 (2008)				
Acceptable Daily Intake: 0-0.015 µg/kg body weight (42 nd JECFA, 1995).				
Residue Definition: Dexamethasone.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	1.0	32 nd (2009)	
Cattle	Liver	2.0	32 nd (2009)	
Cattle	Kidney	1.0	32 nd (2009)	
Cattle	Milk (µg/l)	0.3	32 nd (2009)	
Pig	Muscle	1.0	32 nd (2009)	
Pig	Liver	2.0	32 nd (2009)	
Pig	Kidney	1.0	32 nd (2009)	
Horses	Muscle	1.0	32 nd (2009)	
Horses	Liver	2.0	32 nd (2009)	
Horses	Kidney	1.0	32 nd (2009)	

DICLAZURIL (antiprotozoal agent)				
JECFA Evaluation: 45 (1995); 50 (1998)				
Acceptable Daily Intake: 0-30 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: Diclazuril.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Poultry	Muscle	500	23 rd (1999)	
Poultry	Liver	3000	23 rd (1999)	
Poultry	Kidney	2000	23 rd (1999)	
Poultry	Fat/Skin	1000	23 rd (1999)	
Rabbit	Muscle	500	23 rd (1999)	
Rabbit	Liver	3000	23 rd (1999)	
Rabbit	Kidney	2000	23 rd (1999)	
Rabbit	Fat	1000	23 rd (1999)	
Sheep	Muscle	500	23 rd (1999)	
Sheep	Liver	3000	23 rd (1999)	
Sheep	Kidney	2000	23 rd (1999)	
Sheep	Fat	1000	23 rd (1999)	

DICYCLANIL (insecticide)				
JECFA Evaluation: 54 (2000); 60 (2003)				
Acceptable Daily Intake: 0-7 µg/kg body weight (54 th JECFA, 2000).				
Residue Definition: Dicyclanil.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Sheep	Muscle	150	28 th (2005)	
Sheep	Liver	125	28 th (2005)	
Sheep	Kidney	125	28 th (2005)	
Sheep	Fat	200	28 th (2005)	

DIHYDROSTREPTOMYCIN/STREPTOMYCIN (antimicrobial agent)				
JECFA Evaluation: 43 (1994); 48 (1997); 52 (1999); 58 (2002)				
Acceptable Daily Intake: 0-50 µg/kg body weight (48 th JECFA, 1997). Group ADI for combined residues of dihydrostreptomycin and streptomycin.				
Residue Definition: Sum of dihydrostreptomycin and streptomycin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	600	24 th (2001)	
Cattle	Liver	600	24 th (2001)	
Cattle	Kidney	1000	24 th (2001)	
Cattle	Fat	600	24 th (2001)	
Cattle	Milk	200	26 th (2003)	
Chicken	Muscle	600	24 th (2001)	
Chicken	Liver	600	24 th (2001)	
Chicken	Kidney	1000	24 th (2001)	
Chicken	Fat	600	24 th (2001)	
Pig	Muscle	600	24 th (2001)	
Pig	Liver	600	24 th (2001)	
Pig	Kidney	1000	24 th (2001)	
Pig	Fat	600	24 th (2001)	
Sheep	Muscle	600	24 th (2001)	
Sheep	Liver	600	24 th (2001)	
Sheep	Kidney	1000	24 th (2001)	
Sheep	Fat	600	24 th (2001)	
Sheep	Milk	200	26 th (2003)	

DIMINAZENE (trypanocide)				
JECFA Evaluation: 34 (1989); 42 (1994)				
Acceptable Daily Intake: 0-100 µg/kg body weight (42 nd JECFA, 1994).				
Residue Definition: Diminazene.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	500	22 nd (1997)	
Cattle	Liver	12000	22 nd (1997)	
Cattle	Kidney	6000	22 nd (1997)	
Cattle	Milk (µg/l)	150	22 nd (1997)	Limit of quantitation of the analytical method.

DORAMECTIN (anthelmintic agent)				
JECFA Evaluation: 45 (1995); 52 (1999); 58 (2002); 62 (2004)				
Acceptable Daily Intake: 0-1 µg/kg body weight (58 th JECFA, 2002).				
Residue Definition: Doramectin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	10	22 nd (1997)	High concentration of residues at the injection site over a 35 day period after subcutaneous or intramuscular administration of the drug at the recommended dose.
Cattle	Liver	100	22 nd (1997)	
Cattle	Kidney	30	22 nd (1997)	
Cattle	Fat	150	22 nd (1997)	High concentration of residues at the injection site over a 35 day period after subcutaneous or intramuscular administration of the drug at the recommended dose
Cattle	Milk	15	29 th (2006)	Depending on the route and/or time of administration the use of doramectin in dairy cows may result in extended withdrawal periods in milk. This may be addressed in national/regional regulatory programmes.
Pig	Muscle	5	24 th (2001)	
Pig	Liver	100	24 th (2001)	
Pig	Kidney	30	24 th (2001)	
Pig	Fat	150	24 th (2001)	

EMAMECTIN BENZOATE (antiparasitic agent)				
JECFA Evaluation: 78 (2013)				
Acceptable Daily Intake: ADI of 0–0.5 µg/kg body weight established by the Joint FAO/WHO Meeting on Pesticide Residues (JMPR) in 2011, based on an overall no-observed-adverse effect level (NOAEL) of 0.25 mg/kg body weight per day for neurotoxicity from 14- and 53-week studies in dogs, supported by an overall NOAEL of 0.25 mg/kg body weight per day from 1- and 2-year studies in rats. An uncertainty factor of 500 was applied to the NOAEL, which includes an additional uncertainty factor of 5 to account for the steep dose–response curve and irreversible histopathological effects in neural tissues at the lowest-observed-adverse-effect level (LOAEL) in dogs, as used by JMPR and confirmed by the current Committee (78 th JECFA, 2013).				
Estimated Dietary Exposure: 11 µg/person per day, which represents approximately 37% of the upper bound of the ADI (78 th JECFA, 2013).				
Residue Definition: Emamectin B1a.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Salmon	Muscle	100	38 th (2015)	
Salmon	Fillet	100	38 th (2015)	Muscle plus skin in natural proportion
Trout	Muscle	100	38 th (2015)	
Trout	Fillet	100	38 th (2015)	Muscle plus skin in natural proportion

EPRINOMECTIN (anthelmintic agent)				
JECFA Evaluation: 50 (1998)				
Acceptable Daily Intake: 0-10 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: Eprinomectin B1a.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	26 th (2003)	
Cattle	Liver	2000	26 th (2003)	
Cattle	Kidney	300	26 th (2003)	
Cattle	Fat	250	26 th (2003)	
Cattle	Milk (µg/l)	20	26 th (2003)	

ERYTHROMYCIN (antimicrobial agent)				
JECFA Evaluation: 66 (2006)				
Acceptable Daily Intake: 0-0.7 µg/kg body weight (66 th JECFA, 2006).				
Residue Definition: Erythromycin A				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Chicken	Muscle	100	31 st (2008)	
Chicken	Liver	100	31 st (2008)	
Chicken	Kidney	100	31 st (2008)	
Chicken	Fat	100	31 st (2008)	The MRL includes skin + fat
Chicken	Eggs	50	31 st (2008)	
Turkey	Muscle	100	31 st (2008)	
Turkey	Liver	100	31 st (2008)	
Turkey	Kidney	100	31 st (2008)	
Turkey	Fat	100	31 st (2008)	The MRL includes skin + fat

ESTRADIOL-17BETA (production aid)				
JECFA Evaluation: 25 (1981); 32 (1987); 52 (1999)				
Acceptable Daily Intake: unnecessary (32 nd JECFA, 1987); 0-0.05 µg/kg body weight (52 nd JECFA, 1999).				
Residue Definition: Estradiol-17beta.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Liver	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Kidney	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Fat	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.

FEBANTEL/FENBENDAZOLE/OXFENDAZOLE (anthelmintic agent)				
JECFA Evaluation: 38 (1991); 45 (1995); 50 (1998)				
Acceptable Daily Intake: 0-7 µg/kg body weight (50 th JECFA, 1998). Group ADI				
Residue Definition: Sum of fenbendazole, oxfendazole and oxfendazole sulphone, expressed as oxfendazole sulphone equivalents.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	23 rd (1999)	
Cattle	Liver	500	23 rd (1999)	
Cattle	Kidney	100	23 rd (1999)	
Cattle	Fat	100	23 rd (1999)	
Cattle	Milk (µg/l)	100	23 rd (1999)	
Goat	Muscle	100	23 rd (1999)	
Goat	Liver	500	23 rd (1999)	
Goat	Kidney	100	23 rd (1999)	
Goat	Fat	100	23 rd (1999)	
Horse	Muscle	100	23 rd (1999)	
Horse	Liver	500	23 rd (1999)	
Horse	Kidney	100	23 rd (1999)	
Horse	Fat	100	23 rd (1999)	
Pig	Muscle	100	23 rd (1999)	
Pig	Liver	500	23 rd (1999)	
Pig	Kidney	100	23 rd (1999)	
Pig	Fat	100	23 rd (1999)	
Sheep	Muscle	100	23 rd (1999)	
Sheep	Liver	500	23 rd (1999)	
Sheep	Kidney	100	23 rd (1999)	
Sheep	Fat	100	23 rd (1999)	
Sheep	Milk (µg/l)	100	23 rd (1999)	

FLUAZURON (insecticide)				
JECFA Evaluation: 48 (1997)				
Acceptable Daily Intake: 0-40 µg/kg body weight (48 th JECFA, 1997).				
Residue Definition: Fluazuron.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	200	23 rd (1999)	
Cattle	Liver	500	23 rd (1999)	
Cattle	Kidney	500	23 rd (1999)	
Cattle	Fat	7000	23 rd (1999)	

FLUBENDAZOLE (anthelmintic agent)				
JECFA Evaluation: 40 (1992)				
Acceptable Daily Intake: 0-12 µg/kg body weight (40 th JECFA, 1992).				
Residue Definition: Flubendazole.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Pig	Muscle	10	21 st (1995)	
Pig	Liver	10	21 st (1995)	
Poultry	Muscle	200	21 st (1995)	
Poultry	Liver	500	21 st (1995)	
Poultry	Eggs	400	21 st (1995)	

FLUMEQUINE (antimicrobial agent)				
JECFA Evaluation: 42 (1994); 48 (1997); 54 (2000); 60 (2002); 62 (2004); 66 (2006)				
Acceptable Daily Intake: 0-30 µg/kg body weight (62 nd JECFA, 2004).				
Residue Definition: Flumequine.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	500	28 th (2005)	
Cattle	Liver	500	28 th (2005)	
Cattle	Kidney	3000	28 th (2005)	
Cattle	Fat	1000	28 th (2005)	
Chicken	Muscle	500	28 th (2005)	
Chicken	Liver	500	28 th (2005)	
Chicken	Kidney	3000	28 th (2005)	
Chicken	Fat	1000	28 th (2005)	
Pig	Muscle	500	28 th (2005)	
Pig	Liver	500	28 th (2005)	
Pig	Kidney	3000	28 th (2005)	
Pig	Fat	1000	28 th (2005)	
Sheep	Muscle	500	28 th (2005)	
Sheep	Liver	500	28 th (2005)	
Sheep	Kidney	3000	28 th (2005)	
Sheep	Fat	1000	28 th (2005)	
Trout	Muscle	500	28 th (2005)	Muscle including normal proportion of skin

GENTAMICIN (antimicrobial agent)				
JECFA Evaluation: 43 (1994); 48 (1997); 50 (1998)				
Acceptable Daily Intake: 0-20 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: Gentamicin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	24 th (2001)	
Cattle	Liver	2000	24 th (2001)	
Cattle	Kidney	5000	24 th (2001)	
Cattle	Fat	100	24 th (2001)	
Cattle	Milk (µg/l)	200	24 th (2001)	
Pig	Muscle	100	24 th (2001)	
Pig	Liver	2000	24 th (2001)	
Pig	Kidney	5000	24 th (2001)	
Pig	Fat	100	24 th (2001)	

IMIDOCARB (antiprotozoal agent)				
JECFA Evaluation: 50 (1998); 60 (2003)				
Acceptable Daily Intake: 0-10 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: Imidocarb.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	300	28 th (2005)	
Cattle	Liver	1500	28 th (2005)	
Cattle	Kidney	2000	28 th (2005)	
Cattle	Fat	50	28 th (2005)	
Cattle	Milk	50	28 th (2005)	

ISOMETAMIDIUM (trypanocide)				
JECFA Evaluation: 34 (1989); 40 (1992)				
Acceptable Daily Intake: 0-100 µg/kg body weight (40 th JECFA, 1992).				
Residue Definition: Isometamidium.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	21 st (1995)	
Cattle	Liver	500	21 st (1995)	
Cattle	Kidney	1000	21 st (1995)	
Cattle	Fat	100	21 st (1995)	
Cattle	Milk (µg/l)	100	21 st (1995)	

IVERMECTIN (anthelmintic agent)				
JECFA Evaluation: 36 (1990); 40 (1992); 54 (2000); 58 (2002)				
Acceptable Daily Intake: 0-1 µg/kg body weight (40 th JECFA, 1992).				
Residue Definition: 22,23-Dihydroivermectin B1a (H2B1a).				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Liver	100	20 th (1993)	
Cattle	Fat	40	20 th (1993)	
Cattle	Milk	10	26 th (2003)	
Pig	Liver	15	20 th (1993)	
Pig	Fat	20	20 th (1993)	
Sheep	Liver	15	20 th (1993)	
Sheep	Fat	20	20 th (1993)	

LEVAMISOLE (anthelmintic agent)				
JECFA Evaluation: 36 (1990); 42 (1994)				
Acceptable Daily Intake: 0-6 µg/kg body weight (42 nd JECFA, 1994).				
Residue Definition: Levamisole.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	10	22 nd (1997)	
Cattle	Liver	100	22 nd (1997)	
Cattle	Kidney	10	22 nd (1997)	
Cattle	Fat	10	22 nd (1997)	
Pig	Muscle	10	22 nd (1997)	
Pig	Liver	100	22 nd (1997)	
Pig	Kidney	10	22 nd (1997)	
Pig	Fat	10	22 nd (1997)	
Poultry	Muscle	10	22 nd (1997)	
Poultry	Liver	100	22 nd (1997)	
Poultry	Kidney	10	22 nd (1997)	
Poultry	Fat	10	22 nd (1997)	
Sheep	Muscle	10	22 nd (1997)	
Sheep	Liver	100	22 nd (1997)	
Sheep	Kidney	10	22 nd (1997)	
Sheep	Fat	10	22 nd (1997)	

LINCOMYCIN (antimicrobial agent)				
JECFA Evaluation: 54 (2000); 58 (2002); 62 (2004)				
Acceptable Daily Intake: 0-30 µg/kg body weight (54 th JECFA, 2000).				
Residue Definition: Lincomycin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Milk	150	26 th (2003)	
Chicken	Muscle	200	26 th (2003)	
Chicken	Liver	500	26 th (2003)	
Chicken	Kidney	500	26 th (2003)	
Chicken	Fat	100	26 th (2003)	Additional MRL for skin with adhering fat of 300 µg/kg.
Pig	Muscle	200	26 th (2003)	
Pig	Liver	500	26 th (2003)	
Pig	Kidney	1500	26 th (2003)	
Pig	Fat	100	26 th (2003)	Additional MRL for skin with adhering fat of 300 µg/kg.

MELENGESTROL ACETATE (production aid)				
JECFA Evaluation: 54 (2000); 58 (2002); 62 (2004); 66 (2006) 70 (2008)				
Acceptable Daily Intake: 0-0.03 µg/kg body weight (54 th JECFA, 2000).				
Residue Definition: Melengestrol acetate.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	1	32 nd (2009)	
Cattle	Liver	10	32 nd (2009)	
Cattle	Kidney	2	32 nd (2009)	
Cattle	Fat	18	32 nd (2009)	

MONENSIN (antimicrobial agent)				
JECFA Evaluation:		70 (2008); 75 (2011)		
Acceptable Daily Intake:		0–10 µg/kg body weight on the basis of a NOAEL of 1.14 mg/kg body weight per day and a safety factor of 100 and rounding to one significant figure (70 th JECFA, 2008).		
Estimated Dietary Exposure:		Using the revised MRL, the theoretical maximum daily intake (TMDI) from the 70 th JECFA was recalculated, resulting in a value of 481 µg/person, which represents 80% of the upper bound of the ADI (75 th JECFA, 2011).		
Residue Definition:		Monensin.		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	10	32 nd (2009)	
Cattle	Liver	20	32 nd (2009)	
Cattle	Kidney	100	35 th (2012)	
Cattle	Fat	100	32 nd (2009)	
Cattle	Milk	2	32 nd (2009)	
Sheep	Muscle	10	32 nd (2009)	
Sheep	Liver	20	32 nd (2009)	
Sheep	Kidney	10	32 nd (2009)	
Sheep	Fat	100	32 nd (2009)	
Goats	Muscle	10	32 nd (2009)	
Goats	Liver	20	32 nd (2009)	
Goats	Kidney	10	32 nd (2009)	
Goats	Fat	100	32 nd (2009)	
Chicken	Muscle	10	32 nd (2009)	
Chicken	Liver	10	32 nd (2009)	
Chicken	Kidney	10	32 nd (2009)	
Chicken	Fat	100	32 nd (2009)	
Turkey	Muscle	10	32 nd (2009)	
Turkey	Liver	10	32 nd (2009)	
Turkey	Kidney	10	32 nd (2009)	
Turkey	Fat	100	32 nd (2009)	
Quail	Muscle	10	32 nd (2009)	
Quail	Liver	10	32 nd (2009)	
Quail	Kidney	10	32 nd (2009)	
Quail	Fat	100	32 nd (2009)	

MONEPANTEL (anthelmintic agent)				
JECFA Evaluation: 75 (2011); 78 (2013)				
Acceptable Daily Intake: 0-20 µg/kg body weight on the basis of a no-observed-adverse-effect level (NOAEL) of 1.8 mg/kg body weight per day considering liver effects in mice, and a safety factor of 100, with rounding to one significant figure (75 th JECFA, 2011).				
Estimated Dietary Exposure: Using the model diet and marker residue to total residue ratios of 1.00 for muscle and 0.66 for fat, liver and kidney, and applying a correction factor of 0.94 to account for the mass difference between monepantel sulfone (the marker residue) and monepantel, the EDI is 446 µg/person per day, which represents approximately 37% of the upper bound of the ADI (78 th JECFA, 2013).				
Residue Definition: Monepantel sulfone, expressed as monepantel.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Sheep	Muscle	500	38 th (2015)	
Sheep	Liver	7000	38 th (2015)	
Sheep	Kidney	1700	38 th (2015)	
Sheep	Fat	13000	38 th (2015)	

MOXIDECTIN (anthelmintic agent)				
JECFA Evaluation: 45 (1995); 47 (1996); 48 (1998); 50 (1998)				
Acceptable Daily Intake: 0-2 µg/kg body weight (45 th JECFA, 1995).				
Residue Definition: Moxidectin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	20	22 nd (1997)	Very high concentration and great variation in the level of residues at the injection site in cattle over a 49 day period after dosing.
Cattle	Liver	100	22 nd (1997)	
Cattle	Kidney	50	22 nd (1997)	
Cattle	Fat	500	22 nd (1997)	
Deer	Muscle	20	23 rd (1999)	
Deer	Liver	100	23 rd (1999)	
Deer	Kidney	50	23 rd (1999)	
Deer	Fat	500	23 rd (1999)	
Sheep	Muscle	50	22 nd (1997)	
Sheep	Liver	100	22 nd (1997)	
Sheep	Kidney	50	22 nd (1997)	
Sheep	Fat	500	22 nd (1997)	

NARASIN (antimicrobial agent)				
JECFA Evaluation: 70 (2008); 75 (2011)				
Acceptable Daily Intake: 0-5 µg/kg body weight on the basis of a NOAEL of 0.5 mg/kg body weight per day and a safety factor of 100 (70 th JECFA, 2008).				
Residue Definition: Narasin A.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	15	35 th (2012)	
Cattle	Liver	50	35 th (2012)	
Cattle	Kidney	15	35 th (2012)	
Cattle	Fat	50	35 th (2012)	
Chicken	Muscle	15	32 nd (2009)	
Chicken	Liver	50	32 nd (2009)	
Chicken	Kidney	15	32 nd (2009)	
Chicken	Fat	50	32 nd (2009)	
Pig	Muscle	15	34 th (2011)	
Pig	Liver	50	34 th (2011)	
Pig	Kidney	15	34 th (2011)	
Pig	Fat	50	34 th (2011)	

NEOMYCIN (antimicrobial agent)				
JECFA Evaluation: 43 (1994); 47 (1996); 52 (1999); 58 (2002); 60 (2003)				
Acceptable Daily Intake: 0-60 µg/kg body weight (47 th JECFA, 1996).				
Residue Definition: Neomycin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	500	23 rd (1999)	
Cattle	Liver	500	28 th (2005)	
Cattle	Kidney	10000	28 th (2005)	
Cattle	Fat	500	23 rd (1999)	
Cattle	Milk	1500	28 th (2005)	
Chicken	Muscle	500	23 rd (1999)	
Chicken	Liver	500	23 rd (1999)	
Chicken	Kidney	10000	23 rd (1999)	
Chicken	Fat	500	23 rd (1999)	
Chicken	Eggs	500	23 rd (1999)	
Duck	Muscle	500	23 rd (1999)	
Duck	Liver	500	23 rd (1999)	
Duck	Kidney	10000	23 rd (1999)	
Duck	Fat	500	23 rd (1999)	
Goat	Muscle	500	23 rd (1999)	
Goat	Liver	500	23 rd (1999)	
Goat	Kidney	10000	23 rd (1999)	
Goat	Fat	500	23 rd (1999)	
Pig	Muscle	500	23 rd (1999)	
Pig	Liver	500	23 rd (1999)	
Pig	Kidney	10000	23 rd (1999)	
Pig	Fat	500	23 rd (1999)	
Sheep	Muscle	500	23 rd (1999)	
Sheep	Liver	500	23 rd (1999)	
Sheep	Kidney	10000	23 rd (1999)	
Sheep	Fat	500	23 rd (1999)	
Turkey	Muscle	500	23 rd (1999)	
Turkey	Liver	500	23 rd (1999)	
Turkey	Kidney	10000	23 rd (1999)	
Turkey	Fat	500	23 rd (1999)	

NICARBAZIN (antiprotozoal agent)				
JECFA Evaluation: 50 (1998)				
Acceptable Daily Intake: 0-400 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: N,N'-bis(4-nitropheyl)urea.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Chicken	Muscle	200	23 rd (1999)	Broilers.
Chicken	Liver	200	23 rd (1999)	Broilers.
Chicken	Kidney	200	23 rd (1999)	Broilers.
Chicken	Fat/Skin	200	23 rd (1999)	Broilers.

PHOXIM (insecticide)				
JECFA Evaluation: 52 (1999); 62 (2004)				
Acceptable Daily Intake: 0-4 µg/kg body weight (52 nd JECFA, 1999).				
Residue Definition: Phoxim				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Goat	Muscle	50	26 th (2003)	
Goat	Liver	50	26 th (2003)	
Goat	Kidney	50	26 th (2003)	
Goat	Fat	400	26 th (2003)	
Pig	Muscle	50	26 th (2003)	
Pig	Liver	50	26 th (2003)	
Pig	Kidney	50	26 th (2003)	
Pig	Fat	400	26 th (2003)	
Sheep	Muscle	50	26 th (2003)	
Sheep	Liver	50	26 th (2003)	
Sheep	Kidney	50	26 th (2003)	
Sheep	Fat	400	26 th (2003)	

PIRLIMYCIN (antimicrobial agent)				
JECFA Evaluation: 62 (2004)				
Acceptable Daily Intake: 0-8 µg/kg bw (62 nd JECFA, 2004).				
Residue Definition: Pirlimycin.				
Species	Tissue	MRLs (µg/kg)	CAC	Note
Cattle	Muscle	100	29 th (2006)	
Cattle	Liver	1000	29 th (2006)	
Cattle	Kidney	400	29 th (2006)	
Cattle	Fat	100	29 th (2006)	
Cattle	Milk	100	29 th (2006)	JECFA evaluated the effect of pirlimycin residues on starter cultures and for this reason recommended an MRL of 100 µg/kg of milk. Codex Members may therefore adapt national/regional MRLs in order to address this technological aspect for trade of fresh liquid milk intended for processing using starter culture.

PORCINE SOMATOTROPIN (production aid)				
JECFA Evaluation: 52 (1999)				
Acceptable Daily Intake: Not Specified (52 nd JECFA, 1999).				
Residue Definition: Not applicable.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Pig	Muscle	not specified	26 th (2003)	
Pig	Liver	not specified	26 th (2003)	
Pig	Kidney	not specified	26 th (2003)	
Pig	Fat	not specified	26 th (2003)	

PROGESTERONE (production aid)				
JECFA Evaluation: 25 (1981); 32 (1987); 52 (1999)				
Acceptable Daily Intake: 0-30 µg/kg body weight (52 nd JECFA, 1999).				
Residue Definition: Progesterone.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	unnecessary	21 st (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health
Cattle	Liver	unnecessary	21 st (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health
Cattle	Kidney	unnecessary	21 st (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health
Cattle	Fat	unnecessary	21 st (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health

RACTOPAMINE (production aid)				
JECFA Evaluation: 40 (1992); 62 (2004); 66 (2006)				
Acceptable Daily Intake: 0-1 µg/kg body weight (66 th JECFA, 2006).				
Residue Definition: Ractopamine.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	10	35 th (2012)	
Cattle	Liver	40	35 th (2012)	
Cattle	Kidney	90	35 th (2012)	
Cattle	Fat	10	35 th (2012)	
Pig	Muscle	10	35 th (2012)	
Pig	Liver	40	35 th (2012)	
Pig	Kidney	90	35 th (2012)	
Pig	Fat	10	35 th (2012)	The MRL includes skin + fat

SARAFLOXACIN (antimicrobial agent)				
JECFA Evaluation: 50 (1998)				
Acceptable Daily Intake: 0-0.3 µg/kg body weight (50 th JECFA, 1998).				
Residue Definition: Sarafloxacin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Chicken	Muscle	10	24 th (2001)	
Chicken	Liver	80	24 th (2001)	
Chicken	Kidney	80	24 th (2001)	
Chicken	Fat	20	24 th (2001)	
Turkey	Muscle	10	24 th (2001)	
Turkey	Liver	80	24 th (2001)	
Turkey	Kidney	80	24 th (2001)	
Turkey	Fat	20	24 th (2001)	

SPECTINOMYCIN (antimicrobial agent)				
JECFA Evaluation: 42 (1994); 50 (1998)				
Acceptable Daily Intake: 0-40 µg/kg body weight (42 nd JECFA, 1994).				
Residue Definition: Spectinomycin.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	500	23 rd (1999)	
Cattle	Liver	2000	23 rd (1999)	
Cattle	Kidney	5000	23 rd (1999)	
Cattle	Fat	2000	23 rd (1999)	
Cattle	Milk (µg/l)	200	23 rd (1999)	
Chicken	Muscle	500	23 rd (1999)	
Chicken	Liver	2000	23 rd (1999)	
Chicken	Kidney	5000	23 rd (1999)	
Chicken	Fat	2000	23 rd (1999)	
Chicken	Eggs	2000	23 rd (1999)	
Pig	Muscle	500	23 rd (1999)	
Pig	Liver	2000	23 rd (1999)	
Pig	Kidney	5000	23 rd (1999)	
Pig	Fat	2000	23 rd (1999)	
Sheep	Muscle	500	23 rd (1999)	
Sheep	Liver	2000	23 rd (1999)	
Sheep	Kidney	5000	23 rd (1999)	
Sheep	Fat	2000	23 rd (1999)	

SPIRAMYCIN (antimicrobial agent)				
JECFA Evaluation: 38 (1991); 43 (1994); 47 (1996); 48 (1997)				
Acceptable Daily Intake: 0-50 µg/kg body weight (43 rd JECFA, 1994).				
Residue Definition: Cattle and chickens, sum of spiramycin and neospiramycin; Pigs, spiramycin equivalents (antimicrobially active residues).				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	200	22 nd (1997)	
Cattle	Liver	600	22 nd (1997)	
Cattle	Kidney	300	22 nd (1997)	
Cattle	Fat	300	22 nd (1997)	
Cattle	Milk (µg/l)	200	22 nd (1997)	
Chicken	Muscle	200	22 nd (1997)	
Chicken	Liver	600	22 nd (1997)	
Chicken	Kidney	800	22 nd (1997)	
Chicken	Fat	300	22 nd (1997)	
Pig	Muscle	200	22 nd (1997)	
Pig	Liver	600	22 nd (1997)	
Pig	Kidney	300	22 nd (1997)	
Pig	Fat	300	22 nd (1997)	

SULFADIMIDINE (antimicrobial agent)				
JECFA Evaluation: 34 (1989); 38 (1991); 42 (1994)				
Acceptable Daily Intake: 0-50 µg/kg body weight (42 nd JECFA, 1994).				
Residue Definition: Sulfadimidine.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Milk (µg/l)	25	21 st (1995)	
Not specified	Muscle	100	21 st (1995)	
Not specified	Liver	100	21 st (1995)	
Not specified	Kidney	100	21 st (1995)	
Not specified	Fat	100	21 st (1995)	

TESTOSTERONE (production aid)				
JECFA Evaluation: 25 (1981); 32 (1987); 52 (1999)				
Acceptable Daily Intake: 0-2 µg/kg body weight (52 nd JECFA, 1999).				
Residue Definition: Testosterone.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Liver	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Kidney	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Fat	unnecessary	21 st (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.

THIABENDAZOLE (anthelmintic agent)				
JECFA Evaluation: 40 (1992); 48 (1997); 58 (2002)				
Acceptable Daily Intake: 0-100 µg/kg body weight (40 th JECFA, 1992).				
Residue Definition: Sum of thiabendazole and 5-hydroxythiabendazole.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Liver	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Kidney	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Fat	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Milk (µg/l)	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Muscle	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Liver	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Kidney	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Fat	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Milk (µg/l)	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Muscle	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Liver	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Kidney	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Fat	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Sheep	Muscle	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.

Sheep	Liver	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Sheep	Kidney	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Sheep	Fat	100	21 st (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.

TILMICOSIN (antimicrobial agent)**JECFA Evaluation:** 47 (1996); 54 (2000); 70 (2008)**Acceptable Daily Intake:** 0-40 µg/kg body weight (47th JECFA, 1996).**Residue Definition:** Tilmicosin.

Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	23 rd (1999)	
Cattle	Liver	1000	23 rd (1999)	
Cattle	Kidney	300	23 rd (1999)	
Cattle	Fat	100	23 rd (1999)	
Chicken	Muscle	150	34 th (2011)	
Chicken	Liver	2400	34 th (2011)	
Chicken	Kidney	600	34 th (2011)	
Chicken	Skin/Fat	250	34 th (2011)	
Pig	Muscle	100	23 rd (1999)	
Pig	Liver	1500	23 rd (1999)	
Pig	Kidney	1000	23 rd (1999)	
Pig	Fat	100	23 rd (1999)	
Sheep	Muscle	100	23 rd (1999)	
Sheep	Liver	1000	23 rd (1999)	
Sheep	Kidney	300	23 rd (1999)	
Sheep	Fat	100	23 rd (1999)	
Turkey	Muscle	100	34 th (2011)	
Turkey	Kidney	1200	34 th (2011)	
Turkey	Liver	1400	34 th (2011)	
Turkey	Skin/Fat	250	34 th (2011)	

TRENBOLONE ACETATE (growth promoter)				
JECFA Evaluation: 26 (1982); 27 (1983); 32 (1987); 34 (1989)				
Acceptable Daily Intake: 0-0.02 µg/kg body weight (34 th JECFA, 1989).				
Residue Definition: Cattle muscle, beta-Trenbolone; Cattle liver, alpha-Trenbolone.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	2	21 st (1995)	
Cattle	Liver	10	21 st (1995)	

TRICHLORFON (Metrifonate) (insecticide)				
JECFA Evaluation: 54 (2000); 60 (2003); 66 (2006)				
Acceptable Daily Intake: 0-2 µg/kg bw (60 th JECFA, 2003)				
Residue Definition: JECFA confirmed the MRL for cows's milk and the guidance levels for muscle, liver, kidney and fat of cattle recommended at the 54 th meeting (WHO TRS 900, 2001)				
Species	Tissue	MRLs (µg/kg)	CAC	Notes
Cattle	Milk	50	29 th (2006)	

TRICLABENDAZOLE (anthelmintic agent)				
JECFA Evaluation: 40 (1992); 66 (2006); 70 (2008)				
Acceptable Daily Intake: 0-3 µg/kg body weight (40 th JECFA, 1993).				
Residue Definition: Ketotriclabnedazole				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	250	32 nd (2009)	
Cattle	Liver	850	32 nd (2009)	
Cattle	Kidney	400	32 nd (2009)	
Cattle	Fat	100	32 nd (2009)	
Sheep	Muscle	200	32 nd (2009)	
Sheep	Liver	300	32 nd (2009)	
Sheep	Kidney	200	32 nd (2009)	
Sheep	Fat	100	32 nd (2009)	

TYLOSIN (antimicrobial agent)				
JECFA Evaluation: 70 (2008)				
Acceptable Daily Intake : 0-30 µg/kg body weight based on a microbiological end-point derived from in vitro MIC susceptibility testing and faecal binding data (MIC _{calc} = 1.698) (70 th JECFA, 2008).				
Residue Definition: Tylosin A.				
Species	Tissue	MRLs (µg/kg)	CAC	Notes
Cattle	Muscle	100	32 nd (2009)	
Cattle	Liver	100	32 nd (2009)	
Cattle	Kidney	100	32 nd (2009)	
Cattle	Fat	100	32 nd (2009)	
Cattle	Milk	100	32 nd (2009)	
Pig	Muscle	100	32 nd (2009)	
Pig	Liver	100	32 nd (2009)	
Pig	Kidney	100	32 nd (2009)	
Pig	Fat	100	32 nd (2009)	
Chicken	Muscle	100	32 nd (2009)	
Chicken	Liver	100	32 nd (2009)	
Chicken	Kidney	100	32 nd (2009)	
Chicken	Fat/Skin	100	32 nd (2009)	
Chicken	Eggs	300	32 nd (2009)	

ZERANOL (growth promoter)				
JECFA Evaluation: 26 (1982); 27 (1983); 32 (1987)				
Acceptable Daily Intake : 0-0.5 µg/kg body weight (32 nd JECFA, 1987).				
Residue Definition: Zeranol.				
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	2	21 st (1995)	
Cattle	Liver	10	21 st (1995)	

RISK MANAGEMENT RECOMMENDATIONS (RMRs) FOR RESIDUES OF VETERINARY DRUGS**CARBADOX** (growth promoter)

JECFA evaluation: 36th (1990) and 60th (2003) JECFA

CAC37 (2014)

Recommended risk management measures

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of carbadox or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of carbadox in food. This can be accomplished by not using carbadox in food producing animals.

CHLORAMPHENICOL (antimicrobial agent)

JECFA evaluation: 12th (1968), 32nd (1987), 42nd (1994) and 62nd (2004) JECFA

CAC37 (2014)

Recommended risk management measures

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of chloramphenicol or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of chloramphenicol in food. This can be accomplished by not using chloramphenicol in food producing animals.

CHLORPROMAZINE (tranquilizing agent)

JECFA evaluation: 38th (1991) JECFA

CAC37 (2014)

Recommended risk management measures

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of chlorpromazine or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of chlorpromazine in food. This can be accomplished by not using chlorpromazine in food producing animals.

DIMETRIDAZOLE (antiprotozoal agent)

JECFA evaluation: 34th (1989) JECFA

CAC38 (2015)

Recommended risk management measures

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of dimetridazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of dimetridazole in food. This can be accomplished by not using dimetridazole in food producing animals.

FURAZOLIDONE (antimicrobial agent)

JECFA evaluation: 40th (1992) JECFA

CAC37 (2014)

Recommended risk management measures

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of furazolidone or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of furazolidone in food. This can be accomplished by not using furazolidone in food producing animals.

IPRONIDAZOLE (antiprotozoal agent)**JECFA evaluation:** 34th (1989) JECFA**CAC38** (2015)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of ipronidazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of ipronidazole in food. This can be accomplished by not using ipronidazole in food producing animals.

MALACHITE GREEN (antifungal and antiprotozoal agent)**JECFA evaluation:** 70th (2008) JECFA**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of malachite green or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of malachite green in food. This can be accomplished by not using malachite green in food producing animals.

METRONIDAZOLE (antiprotozoal agent)**JECFA evaluation:** 34th (1989) JECFA**CAC38** (2015)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of metronidazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of metronidazole in food. This can be accomplished by not using metronidazole in food producing animals.

NITROFURAL (antimicrobial agent)**JECFA evaluation:** 40th (1992) JECFA**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of nitrofurantoin or its metabolites¹ in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of nitrofurantoin in food. This can be accomplished by not using nitrofurantoin in food producing animals.

¹ Semicarbazide is not a unique indicator of nitrofurantoin use and low levels can be associated with other legitimate sources.

OLAQUINDOX (antibacterial agent)**JECFA evaluation:** 36th (1990) and 42nd (1994) JECFA**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of olaquinox or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of olaquinox in food. This can be accomplished by not using olaquinox in food producing animals.

RONIDAZOLE (antiprotozoal agent)

JECFA evaluation: 34th (1989) and 42nd (1994) JECFA

CAC38 (2015)

Recommended risk management measures

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of ronidazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of ronidazole in food. This can be accomplished by not using ronidazole in food producing animals.

STILBENES (growth promoter)

JECFA evaluation: 5th (1960) JECFA

IARC evaluation: monograph 100A (2012)

CAC37 (2014)

Recommended risk management measures

In view of the available scientific information, there is no safe level of residues of stilbenes or their metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of stilbenes in food. This can be accomplished by not using stilbenes in food producing animals.